

Census of Antarctic Marine Life
SCAR-Marine Biodiversity Information Network

BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

▶ **CHAPTER 11. THE DYNAMIC BIOGEOGRAPHIC ATLAS PROJECT.**

Danis B., De Broyer C., P. Koubbi P., Van de Putte A.P., 2014.

In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d'Acoz C. d', et al. (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. 466-467.

EDITED BY:

Claude DE BROYER & Philippe KOUBBI (chief editors)

with Huw GRIFFITHS, Ben RAYMOND, Cédric d'UDEKEM
d'ACUZ, Anton VAN DE PUTTE, Bruno DANIS, Bruno DAVID,
Susie GRANT, Julian GUTT, Christoph HELD, Graham HOSIE,
Falk HUETTMANN, Alexandra POST & Yan ROPERT-COUDERT



SCIENTIFIC COMMITTEE ON ANTARCTIC RESEARCH

THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

The “Biogeographic Atlas of the Southern Ocean” is a legacy of the International Polar Year 2007-2009 (www.ipy.org) and of the Census of Marine Life 2000-2010 (www.coml.org), contributed by the Census of Antarctic Marine Life (www.caml.aq) and the SCAR Marine Biodiversity Information Network (www.scarmarbin.be; www.biodiversity.aq).

The “Biogeographic Atlas” is a contribution to the SCAR programmes Ant-ECO (State of the Antarctic Ecosystem) and AnT-ERA (Antarctic Thresholds- Ecosystem Resilience and Adaptation) (www.scar.org/science-themes/ecosystems).

Edited by:

Claude De Broyer (Royal Belgian Institute of Natural Sciences, Brussels)
Philippe Koubbi (Université Pierre et Marie Curie, Paris)
Huw Griffiths (British Antarctic Survey, Cambridge)
Ben Raymond (Australian Antarctic Division, Hobart)
Cédric d’Udekem d’Acoz (Royal Belgian Institute of Natural Sciences, Brussels)
Anton Van de Putte (Royal Belgian Institute of Natural Sciences, Brussels)
Bruno Danis (Université Libre de Bruxelles, Brussels)
Bruno David (Université de Bourgogne, Dijon)
Susie Grant (British Antarctic Survey, Cambridge)
Julian Gutt (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven)
Christoph Held (Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bremerhaven)
Graham Hosie (Australian Antarctic Division, Hobart)
Falk Huettmann (University of Alaska, Fairbanks)
Alix Post (Geoscience Australia, Canberra)
Yan Ropert-Coudert (Institut Pluridisciplinaire Hubert Currien, Strasbourg)

Published by:

The Scientific Committee on Antarctic Research, Scott Polar Research Institute, Lensfield Road, Cambridge, CB2 1ER, United Kingdom (www.scar.org).

Publication funded by:

- The Census of Antarctic Marine Life (Albert P. Sloan Foundation, New York)
- The TOTAL Foundation, Paris.

The “Biogeographic Atlas of the Southern Ocean” shared the *Cosmos Prize* awarded to the Census of Marine Life by the International Osaka Expo’90 Commemorative Foundation, Tokyo, Japan.

Publication supported by:

- The Belgian Science Policy (Belspo), through the Belgian Scientific Research Programme on the Antarctic and the “biodiversity.aq” network (SCAR-MarBIN/ANTABIF)
- The Royal Belgian Institute of Natural Sciences (RBINS), Brussels, Belgium
- The British Antarctic Survey (BAS), Cambridge, United Kingdom
- The Université Pierre et Marie Curie (UPMC), Paris, France
- The Australian Antarctic Division, Hobart, Australia
- The Scientific Steering Committee of CAML, Michael Stoddart (CAML Administrator) and Victoria Wadley (CAML Project Manager)

Mapping coordination and design: Huw Griffiths (BAS, Cambridge) & Anton Van de Putte (RBINS, Brussels)

Editorial assistance: Henri Robert, Xavier Loréa, Charlotte Havermans, Nicole Moortgat (RBINS, Brussels)

Printed by: Altitude Design, Rue Saint Josse, 15, B-1210, Belgium (www.altitude-design.be)

Lay out: Sigrid Camus & Amélie Blaton (Altitude Design, Brussels).

Cover design: Amélie Blaton (Altitude Design, Brussels) and the Editorial Team.

Cover pictures: amphipod crustacean (*Epimeria rubriques* De Broyer & Klages, 1991), image © T. Riehl, University of Hamburg; krill (*Euphausia superba* Dana, 1850), image © V. Siegel, Institute of Sea Fisheries, Hamburg; fish (*Chaenocephalus* sp.), image © C. d’Udekem d’Acoz, RBINS; emperor penguin (*Aptenodytes forsteri* G.R. Gray, 1844), image © C. d’Udekem d’Acoz, RBINS; Humpback whale (*Megaptera novaeangliae* (Borowski, 1781)), image © L. Kindermann, AWI.

Online dynamic version :

A dynamic online version of the Biogeographic Atlas is available on the SCAR-MarBIN / AntaBIF portal : atlas.biodiversity.aq.

Recommended citation:

For the volume:

De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d’Acoz C. d’, Van de Putte A.P., Danis B., David B., Grant S., Gutt J., Held C., Hosie G., Huettmann F., Post A., Ropert-Coudert Y. (eds.), 2014. Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, XII + 498 pp.

For individual chapter:

(e.g.) Crame A., 2014. Chapter 3.1. Evolutionary Setting. In: De Broyer C., Koubbi P., Griffiths H.J., Raymond B., Udekem d’Acoz C. d’, *et al.* (eds.). Biogeographic Atlas of the Southern Ocean. Scientific Committee on Antarctic Research, Cambridge, pp. xx-yy.

ISBN: 978-0-948277-28-3.



This publication is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License

11. Going dynamic: a vision for the Biogeographic Atlas of the Southern Ocean (BASO)

Bruno Danis¹, Claude De Broyer², Philippe Koubbi³, Anton P. Van de Putte²

¹ Marine Biology Laboratory, Université Libre de Bruxelles, Brussels, Belgium

² Royal Belgian Institute of Natural Sciences, Brussels, Belgium

³ Sorbonne Universités. UMR BOREA 7208, Université Pierre et Marie Curie, Muséum national d'Histoire naturelle, Paris, France

1. Context

Biogeographic information is of primary importance for monitoring biodiversity and detecting impacts of environmental changes, developing the ecoregionalisation of the Southern Ocean, designing marine protected areas, conducting comparative biogeographic analyses, and discovering marine biodiversity hotspots. Stressed by fast-paced change, and strong environmental pressure, the importance of having high quality data openly available online has become part of the Antarctic researchers culture. As a result, more and more biodiversity primary data is made available through dedicated information systems, which take the form of web or data portals, offering free and open access to highly needed information.

The multi-authored «Biogeographic Atlas of the Southern Ocean» (BASO), builds upon information networks and their collaborative communities, to provide an up-to-date synthesis of the Antarctic biogeographic knowledge. The first version of the BASO is taking the form of a «classic» printout version. To take full advantage of the existing informatics tools, in particular biodiversity informatics and web tools, a dynamic approach has been envisaged from the beginning of the project to follow-up on the initial, static version. This dynamic version will give birth to a new interoperable system, building customized biogeographic synthesis pages on-the-fly, using the content of relevant information systems as it improves over time.

It is important to stress that only by ensuring community-driven developments, scientific independence, and high levels of data interoperability will this initiative meet the success it deserves and involve a wide range of stakeholders.

2. Overall scope of BASO

Adopting the same approach, the dynamic, digital version will be made available through a dedicated portal. This dynamic BASO will build its content from a blend of authoritative sources, including of course data assembled in the framework of the static version, but also taking advantage of a dynamic approach, allowing re-generation of maps including the latest available taxonomic and biogeographic data from various sources. The dynamic BASO could also include advanced functionalities, such as the capacity to build tailor-made maps, the possibility to tweak the modelisation parameters, the inclusion of data available from different databases (Register of Antarctic Marine Species (RAMS), SCAR-MarBIN, ANTABIF, Antarctic Field Guides, GenBank,...), allowing the end-users to provide feedback on data or models, eventually turning the BASO into a community-driven platform entailed to publish and share maps generated using the most accurate and up-to-date information.

Hence, the dynamic version will complete and expand the hard copy version, and offer in addition – using appropriate technical solution such as webservice and APIs (Application Programming Interfaces) - visualisation tools as well as environmental/model datasets in appropriate formats directly useful for Geographical Information Systems (GIS) users. This approach would be particularly useful in the framework of habitat modelling and prediction of potential climate-related changes. Habitat modelling allows predicting species potential distribution according to the species relation to a set of environmental variables. The species modelling is not possible for all of the considered species, but the identification of areas of major biodiversity change is nevertheless achievable. Two workshops identified in 2010 and 2011 the methodology that will be used for the Atlas. (see Koubbi *et al.* 2010, 2011; Loots *et al.* 2007; Pinkerton *et al.* 2010).

In designing the dynamic version of BASO, special care should be devoted to ease the access to data, documenting the content (using metadata from the Global Change Master Directory, SERF format, or GBIF Ecological Markup Language (EML)) and using widely accepted Geospatial standards (Open Geospatial Consortium (OGC)) to ensure interoperability and direct usability of the data in as many contexts as possible.

To the best extent feasible, emerging datatypes (tracking, genetics, microbial, ROV imagery,...) should be anticipated in the design of the data and workflows for the BASO.

3. Stakeholders

A dynamic version of the BASO will probably attract a wide variety of users from various horizons, but will also be useful for connected initiatives which will integrate the relevant data and workflows generated in the framework of the BASO, in a multidirectional fashion.

Regarding the users, there are high chances that the primary end-users for the BASO data portal will be the Antarctic researchers (biogeographers, taxonomists, ecologists, molecular biologists, conservationists, modellers,...).

However, BASO typically represents a cross-disciplinary exercise. This is especially relevant in the context of the new SCAR research programmes, which have been specifically designed to promote cross-linked approaches. Other stakeholders include conservationists, policy makers (Antarctic Treaty System: Committee on Environmental Protection (CEP), Commission for the Conservation of the Antarctic Marine Living Resources (CCAMLR),...), as well as students, the education & outreach community or even the general public. Networked initiatives such as ANTABIF, OBIS, GBIF, SOOS, the Antarctic Environments Portal will also be incorporated in the multidirectional data flows.

4. Requirements

Even if it is still early to anticipate all the fine-scale requirements and functionalities of the dynamic version of BASO at this stage, a set of general considerations can nevertheless be proposed. BASO should be considered as a tool to facilitate discovery, generation and access to both factual and predictive biogeographic data, including the possibility to offer access to additional sources of information at various levels of biological complexity.

As a development platform, BASO will be open to both data end-users and data publishers. Future managers of the BASO are expected to be aware of the needs of data end-users and the wishes and needs of data publishers. To the best extent possible, the wishes of the relevant communities regarding BASO functionalities should be expressed in the flexibility and scalability of the tool itself. Based on an iterative developmental process, feedback can be expected from the users community in order to match their needs and the functionalities of the BASO.




Figure 1 Wireframe describing the general concept for the Dynamic version of the Biogeographic Atlas content building, and associated workflows

A proposed general workflow to generate new predictive maps is described in Fig 1. New occurrence data is published and documented by data providers through the ANTABIF Integrated Publishing Toolkit (IPT). New data is exposed to ANTABIF, GBIF and OBIS for harvesting, and also to the BASO R server, which processes the data and incorporates it in newly generated predictive maps. The maps are published using the ANTABIF geoserver, and made available and searchable through the BASO webportal.

A set of basic requirements, which should be met during the development of the dynamic version of the BASO are described below:

Data consistency between BASO instances and ANTABIF. The BASO will include the possibility to acquire contextual data from the ANTABIF central index. The process of synchronizing data between BASO and ANTABIF should be more or less automated, depending of the type of data to be synchronised:

1. occurrence data - format Darwin Core (DwC) table, all data kept by ANTABIF, extending to all fields available, as defined on the relevant GBIF resource webpage.
2. species lists - format DwC table for species lists extension, all data kept by ANTABIF, not only the indexed data.

3. metadata - in GBIF EML and GCMD DIF formats, again not only indexed data but resource-level metadata.
4. maps - generated from the BASO R-server are posted on the ANTABIF geoserver using OGC-compliant standards.
5. multimedia metadata - according to the MRTG schema for the multimedia metadata format, all data, not only the index.

Free & Open source solutions. To the best extent possible, the BASO system should be based on layers of open source solutions such as PHP, MySQL, Apache. Open source application development frameworks or libraries such as JQuery (javascript), CakePHP, that streamline and simplify both code development and updates should also be used. Web standards (HTML5, CSS3, JQuery) should also be considered. Easy maintenance of the application is an important criteria.

Compliant with the ANTABIF Informatics Infrastructure / Architecture. To ensure the interoperability of the BASO with other ANTABIF tools such as IPT, AFG, ..., BASO should be developed in a manner that is compliant with ANTABIF informatics infrastructure and architecture. The interoperability of the BASO with other ANTABIF informatics tools should be aiming at optimizing current development, and ensuring the same standards are adopted and that resulting data flows are kept simple and consistent (i.e. not affecting the data quality of the data system as a whole).

Robust. The BASO system should be thoroughly tested to ensure robustness on two fronts: (1) the BASO should be able to cope (without crashing) with a wide range of stress situations, and it should perform reasonably under important workload (queries and data discovery). User experience should be as intuitive and friendly as possible; any errors should be trapped and not seen by users, but systematically addressed as priority requirements or issues; (2) it should also allow the management of dependencies on external services (for example, external data calls).

Scalable. Scalability is an essential feature, as the BASO should be able to cope with rapidly increasing data masses at least equivalent to that available from the current ANTABIF content, in a fashion that does not impact the end-user's experience (delays). The BASO should be adaptive, in order to allow the inclusion of major data loads in the future.

Modular. The BASO should be built in a modular fashion, which ensures that component functionalities can be adjusted, added, improved, removed or exchanged without impacting the application as a whole.

Extensible. The BASO development and implementation should follow design principles that will enable future growth, both in terms of the addition of new functionality or through modification of existing functionalities. In addition, BASO should provide the possibility to consume services and functions that are created elsewhere. Examples might include taxonomic or biogeographic data sources, mapping functions or web map/feature services.

Intuitive and friendly end-user interface. The interface for end-users should be intuitive and easy to use, without requiring extensive technical assistance, support or off line user guides.

Documentation and code maintenance. A clear and standard practice for the maintenance and documentation of code should be set in place, agreed and communicated as a best practice.

Open collaborative development. The development of the application should be consistent with open source, allowing the core code and any modules developed to be improved over time.

Community-driven. By including opportunities for user feedback, the system should evolve by iterations and respond to user needs, with a focus on the SCAR users community. User interactions can also happen by using stored queries. The implication of the community is a core requirement for the

BASO, and the objective is to ensure that the tool has enough buy-in for it to be allowed to evolve in the future.

Discoverable through search engines. The BASO should have appropriate html metadata tags that allow the tool itself to be discovered through search engines. Metadata made available to search engines should aim primarily at making the BASO data contents easily discoverable through e.g. Google.

Availability of APIs and Web Services. The design and creation of Application Programming Interfaces (APIs) that expose BASO functions should be explored to allow third parties to use BASO functionalities. Web services should also be explored.

6. Perspectives

The Biogeographic Atlas of the Southern Ocean (BASO) is an unprecedented effort to reconcile vast amounts of knowledge on a single interface. This interface is first envisioned as a hard copy, with the potential to become a dynamic, online information system.

Built in a modular fashion, the dynamic BASO should be able to cope with fast-evolving data flows, biodiversity informatics technologies and user needs. Constantly up-to-date, the system could not only play an important role for research purposes (representing a powerful integration tool) but also for conservationists or policy makers. Emerging initiatives, such as the Antarctic Environments Portal (www.environments.aq), or the Southern Ocean Observing System (www.soos.aq), will immediately benefit from the modelling power of the BASO workflows.

In addition, these workflows can be further shared with other integrative infrastructures, such as the Lifewatch initiative (www.lifewatch.eu).

A hundred and forty contributors have worked on the static version of the project so far, and will be directly or indirectly associated to the dynamic version of the BASO. There are high hopes that this interest and commitment will grow over time, and that other communities will join the project, and further broaden its scope. During IPY, the polar data science community was humbled by how the rapid and dramatic changes in the polar regions increased the expectations for an efficient access to rigorous, scrupulously documented data. A next priority for the BASO community is to identify mechanisms for collaborative international funding that supports the necessarily global aspects of such interdisciplinary projects. The BASO addresses deeply interdisciplinary problems. As such, as a key component of the polar scientific data ecosystem, it requires continued organic adaptation by technology, people, organizations, and society.

The latest information about the developments of the dynamic version of BASO can be found on the following url: <http://atlas.biodiversity.aq>

References

- Koubbi, P., Moteki, M., Duhamel, G., Goarant, A., Hulley, P.-A., O'Driscoll, R., Ishimaru, T., *et al.*, 2011. Ecoregionalization of myctophid fish in the Indian sector of the Southern Ocean: Results from generalized dissimilarity models. *Deep Sea Research Part II: Topical Studies in Oceanography*, **58(1-2)**, 170–180. doi:10.1016/j.dsr2.2010.09.007
- Koubbi, P., Ozouf-Costaz, C., Goarant, A., Moteki, M., Hulley, P.-A., Causse, R., Dettai, A., *et al.*, 2010. Estimating the biodiversity of the East Antarctic shelf and oceanic zone for ecoregionalisation: Example of the ichthyofauna of the CEAMARC (Collaborative East Antarctic Marine Census) CAML surveys. *Polar Science*, **4(2)**, 18. doi:10.1016/j.polar.2010.04.012
- Loots, C., Koubbi, P., Duhamel, G., 2007. Habitat modelling of *Electrona antarctica* (Myctophidae, Pisces) in Kerguelen by generalized additive models and geographic information systems. *Polar Biology*, **30**, 951–959.
- Pinkerton, M.H., Smith, A.N.H., Raymond, B., Hosie, G.W., Sharp, B., Leathwick, J.R., Bradford-Grieve, J.M., 2010. Spatial and seasonal distribution of adult *Oithona similis* in the Southern Ocean: Predictions using boosted regression trees. *Deep Sea Research I*, **57(4)**, 469–485. doi:10.1016/j.dsr.2009.12.010

THE BIOGEOGRAPHIC ATLAS OF THE SOUTHERN OCEAN

Scope

Biogeographic information is of fundamental importance for discovering marine biodiversity hotspots, detecting and understanding impacts of environmental changes, predicting future distributions, monitoring biodiversity, or supporting conservation and sustainable management strategies.

The recent extensive exploration and assessment of biodiversity by the Census of Antarctic Marine Life (CAML), and the intense compilation and validation efforts of Southern Ocean biogeographic data by the SCAR Marine Biodiversity Information Network (SCAR-MarBIN / OBIS) provided a unique opportunity to assess and synthesise the current knowledge on Southern Ocean biogeography.

The scope of the Biogeographic Atlas of the Southern Ocean is to present a concise synopsis of the present state of knowledge of the distributional patterns of the major benthic and pelagic taxa and of the key communities, in the light of biotic and abiotic factors operating within an evolutionary framework. Each chapter has been written by the most pertinent experts in their field, relying on vastly improved occurrence datasets from recent decades, as well as on new insights provided by molecular and phylogeographic approaches, and new methods of analysis, visualisation, modelling and prediction of biogeographic distributions.

A dynamic online version of the Biogeographic Atlas will be hosted on www.biodiversity.aq.

The Census of Antarctic Marine Life (CAML)

CAML (www.caml.aq) was a 5-year project that aimed at assessing the nature, distribution and abundance of all living organisms of the Southern Ocean. In this time of environmental change, CAML provided a comprehensive baseline information on the Antarctic marine biodiversity as a sound benchmark against which future change can reliably be assessed. CAML was initiated in 2005 as the regional Antarctic project of the worldwide programme Census of Marine Life (2000-2010) and was the most important biology project of the International Polar Year 2007-2009.

The SCAR Marine Biodiversity Information Network (SCAR-MarBIN)

In close connection with CAML, SCAR-MarBIN (www.scarmarbin.be, integrated into www.biodiversity.aq) compiled and managed the historic, current and new information (i.a. generated by CAML) on Antarctic marine biodiversity by establishing and supporting a distributed system of interoperable databases, forming the Antarctic regional node of the Ocean Biogeographic Information System (OBIS, www.iobis.org), under the aegis of SCAR (Scientific Committee on Antarctic Research, www.scar.org). SCAR-MarBIN established a comprehensive register of Antarctic marine species and, with biodiversity.aq provided free access to more than 2.9 million Antarctic georeferenced biodiversity data, which allowed more than 60 million downloads.

The Editorial Team



Claude DE BROYER is a marine biologist at the Royal Belgian Institute of Natural Sciences in Brussels. His research interests cover structural and ecofunctional biodiversity and biogeography of crustaceans, and polar and deep sea benthic ecology. Active promoter of CAML and ANDEEP, he is the initiator of the SCAR Marine Biodiversity Information Network (SCAR-MarBIN). He took part to 19 polar expeditions.




Huw GRIFFITHS is a marine Biogeographer at the British Antarctic Survey. He created and manages SOMBASE, the Southern Ocean Mollusc Database. His interests include large-scale biogeographic and ecological patterns in space and time. His focus has been on molluscs, bryozoans, sponges and pycnogonids as model groups to investigate trends at high southern latitudes.



Cédric d'UDEKEM d'ACOZ is a research scientist at the Royal Belgian Institute of Natural Sciences, Brussels. His main research interests are systematics of amphipod crustaceans, especially of polar species and taxonomy of decapod crustaceans. He took part to 2 scientific expeditions to Antarctica on board of the *Polarstern* and to several sampling campaigns in Norway and Svalbard.



Bruno DANIS is an Associate Professor at the Université Libre de Bruxelles, where his research focuses on polar biodiversity. Former coordinator of the www.scarmarbin.be and antabif.be projects, he is a leading member of several international committees, such as OBIS or the SCAR Expert Group on Antarctic Biodiversity Informatics. He has published papers in various fields, including ecotoxicology, physiology, biodiversity informatics, polar biodiversity or information science.



Susie GRANT is a marine biogeographer at the British Antarctic Survey. Her work is focused on the design and implementation of marine protected areas, particularly through the use of biogeographic information in systematic conservation planning.



Christoph HELD is a Senior Research Scientist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven. He is a specialist in molecular systematics and phylogeography of Antarctic crustaceans, especially isopods.



Falk HUETTMANN is a 'digital naturalist' he works on three poles (Arctic, Antarctic and Hindu-Kush Himalaya) and elsewhere (marine, terrestrial and atmosphere). He is based with the university of Alaska-Fairbank (UAF) and focuses primarily on effective conservation questions engaging predictions and open access data.



Philippe KOUUBI is professor at the University Pierre et Marie Curie (Paris, France) and a specialist in Antarctic fish ecology and biogeography. He is the Principal Investigator of projects supported by IPEV, the French Polar Institute. As a French representative to the CCAMLR Scientific Committee, his main input is on the proposal of Marine Protected Areas. His other field of research is on the ecoregionalisation of the high seas.



Ben RAYMOND is a computational ecologist and exploratory data analyst, working across a variety of Southern Ocean, Antarctic, and wider research projects. His areas of interest include ecosystem modelling, regionalisation and marine protected area selection, risk assessment, animal tracking, seabird ecology, complex systems, and remote sensed data analyses.



Anton VAN DE PUTTE works at the Royal Belgian Institute for Natural Sciences (Brussels, Belgium). He is an expert in the ecology and evolution of Antarctic fish and is currently the Science Officer for the Antarctic Biodiversity Portal www.biodiversity.aq. This portal provides free and open access to Antarctic Marine and terrestrial biodiversity of the Antarctic and the Southern Ocean.



Bruno DAVID is CNRS director of research at the laboratory BIOGÉOSCIENCES, University of Burgundy. His works focus on evolution of living forms, with and more specifically on sea urchins. He authored a book and edited an extensive database on Antarctic echinoids. He is currently President of the scientific council of the Muséum National d'Histoire Naturelle (Paris), and Deputy Director at the CNRS Institute for Ecology and Environment.




Julian GUTT is a marine ecologist at the Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, and professor at the Oldenburg University, Germany. He participated in 13 scientific expeditions to the Antarctic and was twice chief scientist on board *Polarstern*. He is member of the SCAR committees ACCE and AN-T-ERA (as chief officer). Main foci of his work are: biodiversity, ecosystem functioning and services, response of marine systems to climate change, non-invasive technologies, and outreach.



Graham HOSIE is Principal Research Scientist in zooplankton ecology at the Australian Antarctic Division. He founded the SCAR Southern Ocean Continuous Plankton Recorder Survey and is the Chief Officer of the SCAR Life Sciences Standing Scientific Group. His research interests include the ecology and biogeography of plankton species and communities, notably their response to environmental changes. He has participated in 17 marine science voyages to Antarctica.



Alexandra POST is a marine geoscientist, with expertise in benthic habitat mapping, sedimentology and geomorphic characterisation of the seafloor. She has worked at Geoscience Australia since 2002, with a primary focus on understanding seafloor processes and habitats on the East Antarctic margin. Most recently she has led work to understand the biophysical environment beneath the Amery Ice Shelf, and to characterise the habitats on the George V Shelf and slope following the successful CAML voyages in that region.



Yan ROPERT COUDERT spent 10 years at the Japanese National Institute of Polar Research, where he graduated as a Doctor in Polar Sciences in 2001. Since 2007, he is a permanent researcher at the CNRS in France and the director of a polar research programme (since 2011) that examines the ecological response of Adélie penguins to environmental changes. He is also the secretary of the Expert Group on Birds and Marine Mammals and of the Life Science Group of the Scientific Committee on Antarctic Research.

